


# Exploring the cosmological dark matter coincidence with infrared fixed points

---

Alex Ritter, Raymond Volkas

arXiv: 2210.11011



# The cosmological coincidence


Large range of DM candidates

- Axions, WIMPs, sterile neutrinos, PBHs...
- How to guide our model building?

**Clues** from current observational evidence:

- Apparent coincidence in the mass densities of visible and dark matter

$$\Omega_{DM} \simeq 5\Omega_{VM}$$



TODAY

[https://wmap.gsfc.nasa.gov/universe/uni\\_matter.html](https://wmap.gsfc.nasa.gov/universe/uni_matter.html)

Parameter	TT,TE,EE+lowE+lensing+BAO 68% limits
$\Omega_b h^2$ . . . . .	$0.02242 \pm 0.00014$
$\Omega_c h^2$ . . . . .	$0.11933 \pm 0.00091$

Planck 2018, arXiv: 1807.06209

# Explaining the coincidence

Our goal is to build models in which the mass densities of visible and dark matter are naturally of a similar order of magnitude

This is a problem in two parts:  $\Omega_X = n_X m_X / \rho_c$


Relating number densities

$$n_B \sim n_D$$

Relating particle masses

$$m_B \sim m_D$$

# Relating number densities - ADM



The **visible** number density: asymmetry between baryons and antibaryons (or a nonzero baryon number  $B_V$ )

$$\Omega_{\text{VM}} \equiv \frac{\rho_p - \rho_{\bar{p}}}{\rho_c} \simeq \frac{\rho_p}{\rho_c}$$

proton  
critical

In Asymmetric Dark Matter models there exists a similar asymmetry in a dark baryon number  $B_D$

Wide range of ADM literature where  $n_B \sim n_D$

Most ADM models do not motivate  $m_B \sim m_D$

These are **not** satisfactory explanations of the coincidence problem

# Relating particle masses



The **visible** baryon mass: largely from the QCD confinement scale  $\Lambda_{QCD}$


**Dark matter:** baryon-like bound states of a QCD-like confining gauge group  $SU(3)_{dQCD}$  with

$$\Lambda_{QCD} \sim \Lambda_{dQCD}$$

There are two main ways to achieve this:

1. Introduce a symmetry between  $SU(3)_{QCD}$  and  $SU(3)_{dQCD}$  e.g. [AR, Volkas: 2101.07421](#)
2. The gauge couplings of the two groups can evolve to some ***infrared fixed point***

# Infrared fixed points & Dark QCD



## Bai and Schwaller [1306.4676]

- Dark QCD –  $SU(3)_{dQCD}$
- New fields
  - All at a heavy mass scale  $M$
  - Except for some light quarks (to be confined into dark baryons)

Field	$SU(3)_{QCD} \times SU(3)_{dQCD}$	Mass	Multiplicity
Fermion	$(\mathbf{3}, \mathbf{1})$	$M$	$n_{f_{c,h}}$
	$(\mathbf{1}, \mathbf{3})$	$< \Lambda_{dQCD}$	$n_{f_{d,l}}$
		$M$	$n_{f_{d,h}}$
	$(\mathbf{3}, \mathbf{3})$	$M$	$n_{f_j}$
Scalar	$(\mathbf{3}, \mathbf{1})$	$M$	$n_{s_c}$
	$(\mathbf{1}, \mathbf{3})$	$M$	$n_{s_d}$
	$(\mathbf{3}, \mathbf{3})$	$M$	$n_{s_j}$

# Infrared fixed points & Dark QCD

## Bai and Schwaller [1306.4676]

- Dark QCD –  $SU(3)_{dQCD}$
- New fields
  - All at a heavy mass scale  $M$
  - Except for some light quarks (to be confined into dark baryons)

Get coupled two-loop beta-functions for the coupling constants

$$\begin{aligned} \beta_c = & \frac{g_c^3}{16\pi^2} \left[ \frac{2}{3} (n_{f_c} + 3n_{f_j}) + \frac{1}{6} (n_{s_c} + 3n_{s_j}) - 11 \right] \\ & + \frac{g_c^5}{(16\pi^2)^2} \left[ \frac{38}{3} (n_{f_c} + 3n_{f_j}) + \frac{11}{3} (n_{s_c} + 3n_{s_j}) - 102 \right] \\ & + \frac{g_c^3 g_d^2}{(16\pi^2)^2} [8n_{f_j} + 8n_{s_j}], \end{aligned}$$

Field	$SU(3)_{QCD} \times SU(3)_{dQCD}$	Mass	Multiplicity
Fermion	$(\mathbf{3}, \mathbf{1})$	$M$	$n_{f_{c,h}}$
	$(\mathbf{1}, \mathbf{3})$	$< \Lambda_{dQCD}$	$n_{f_{d,l}}$
	$(\mathbf{3}, \mathbf{3})$	$M$	$n_{f_{d,h}}$
Scalar	$(\mathbf{3}, \mathbf{1})$	$M$	$n_{s_c}$
	$(\mathbf{1}, \mathbf{3})$	$M$	$n_{s_d}$
	$(\mathbf{3}, \mathbf{3})$	$M$	$n_{s_j}$

# Infrared fixed points & Dark QCD

## Bai and Schwaller [1306.4676]

- Dark QCD –  $SU(3)_{dQCD}$
- New fields
  - All at a heavy mass scale  $M$
  - Except for some light quarks (to be confined into dark baryons)

Get coupled two-loop beta-functions for the coupling constants

$$\begin{aligned} \beta_c = & \frac{g_c^3}{16\pi^2} \left[ \frac{2}{3} (n_{f_c} + 3n_{f_j}) + \frac{1}{6} (n_{s_c} + 3n_{s_j}) - 11 \right] \\ & + \frac{g_c^5}{(16\pi^2)^2} \left[ \frac{38}{3} (n_{f_c} + 3n_{f_j}) + \frac{11}{3} (n_{s_c} + 3n_{s_j}) - 102 \right] \\ & + \frac{g_c^3 g_d^2}{(16\pi^2)^2} [8n_{f_j} + 8n_{s_j}], \end{aligned}$$


Field	$SU(3)_{QCD} \times SU(3)_{dQCD}$	Mass	Multiplicity
Fermion	$(\mathbf{3}, \mathbf{1})$	$M$	$n_{f_{c,h}}$
	$(\mathbf{1}, \mathbf{3})$	$< \Lambda_{dQCD}$	$n_{f_{d,l}}$
	$(\mathbf{3}, \mathbf{3})$	$M$	$n_{f_{d,h}}$
Scalar	$(\mathbf{3}, \mathbf{1})$	$M$	$n_{s_c}$
	$(\mathbf{1}, \mathbf{3})$	$M$	$n_{s_d}$
	$(\mathbf{3}, \mathbf{3})$	$M$	$n_{s_j}$

Depending on the field content (model), can have an **infrared fixed point** where

$$\beta_c(\alpha_s^*, \alpha_d^*) = \beta_d(\alpha_s^*, \alpha_d^*) = 0$$



# Relating the confinement scales




1. the coupling constants evolve to the fixed point  $(\alpha_s^*, \alpha_d^*)$  regardless of their initial value in the UV
2. The decoupling scale  $M$  is determined by matching the running of  $\alpha_s$  below  $M$  with experiment
3. The dark confinement scale  $\Lambda_{\text{dQCD}}$  is then determined by running  $\alpha_d$  until it reaches  $\pi/4$

Process: **model (field content)**  $\rightarrow \{\alpha_s^*, \alpha_d^*\} \rightarrow M \rightarrow \Lambda_{\text{dQCD}}$

# Initial UV conditions

**However!**

Couplings do not always evolve to their IRFP values by the decoupling scale




New process: **model,  $\{\alpha_s^{UV}, \alpha_d^{UV}\} \rightarrow M \rightarrow \Lambda_{dQCD}$**

1. ~~the coupling constants evolve to the fixed point  $(\alpha_s^*, \alpha_d^*)$  regardless of their initial value in the UV~~
2. The decoupling scale  $M$  is determined by matching the running of  $\alpha_s$  below  $M$  with experiment
3. The dark confinement scale  $\Lambda_{dQCD}$  is then determined by running  $\alpha_d$  until it reaches  $\pi/4$

# Explaining the coincidence

**Goal** : obtain similar confinement scales  
for visible and dark QCD

$$0.2 \text{ GeV} < \Lambda_{dQCD} < 5 \text{ GeV}$$




# Explaining the coincidence

**Goal** : obtain similar confinement scales for visible and dark QCD

$$0.2 \text{ GeV} < \Lambda_{dQCD} < 5 \text{ GeV}$$

Define  $\varepsilon_\nu$



- ‘viable fraction’ of  $\{\alpha_s^{UV}, \alpha_d^{UV}\}$  parameter space
- simple heuristic for the naturalness of a given model



# Results

First looked at models with at most 3 of each new field



- **12,288** models
- **155** with a perturbative IRFP





# Results

First looked at models with at most 3 of each new field

- **12,288** models
- **155** with a perturbative IRFP







### Issue:

$M < 1 \text{ TeV}$  for much of the viable parameter space

New sub-TeV coloured fields would be produced at colliders





# Looking for models with large M

Look at models with large  $n_{s_j}$  (# of joint scalars)

- Increases the magnitude of all beta-function coefficients

**188** models with  $n_{s_j} > 10$  and one-loop beta-function coefficients between -0.1 and 0




# Conclusions

Cosmological coincidence inspires interesting model building


We've found a set of phenomenologically viable models that could naturally have  $m_B \sim m_D$

Questions?



# Backup Slides

---




# Why is it a coincidence?

Unrelated mechanisms generate the mass density of visible baryons and most dark matter candidates

- **Visible baryons:** baryon-antibaryon asymmetry from baryogenesis
- **WIMPs:** thermal freeze-out
- **Axions:** misalignment mechanism

*A priori* we would not expect the dark and visible mass densities to be on the same order of magnitude




Stephen J. Lonsdale, Thesis (2018)

# Threshold corrections

Heavy fields can still affect beta-functions at energies below their mass scale  $M$

Need to apply threshold conditions at a decoupling scale  $\mu_0 = \mathcal{O}(M)$



This introduces an uncertainty into  $M$ ,  $\Lambda_{dQCD}$  for a given  $\{\alpha_s^{UV}, \alpha_d^{UV}\}$




# Asymptotic Freedom

With coupled beta-functions, asymptotic freedom depends on the values of the gauge couplings

We only work with couplings that are perturbative below the Planck scale, so do consider non-asymptotically free regions



# Bai-Schwaller results



Model	$n_{f_c}$	$n_{f_d}$	$n_{f_j}$	$n_{s_c}$	$n_{s_d}$	$n_{s_j}$	$\alpha_s^*$	$\alpha_d^*$
A	6	5	3	0	2	0	0.095	0.175
B	6	6	3	1	0	0	0.083	0.120
C	6	6	3	2	2	0	0.070	0.070
D	7	7	2	2	0	2	0.078	0.168
E	7	7	2	2	1	2	0.090	0.133
F	8	8	2	2	0	1	0.074	0.149
G	8	8	2	2	1	1	0.082	0.118